


LECTURE 10 – Kronig Penney Model

1.12 Kronig-Penney Model

Consider the following idealized crystal potential:

We assume $E < V_0$.



for $0 \leq x \leq a$:

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2} E\psi = 0$$

and

$$\psi = A \sin \alpha x + B \cos \alpha x \quad \frac{d\psi}{dx} = A\alpha \cos \alpha x - B\alpha \sin \alpha x \quad \alpha = \frac{\sqrt{2mE}}{\hbar}$$

and for $-b \leq x \leq 0$:

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2} (E - V_0)\psi = 0$$

Solution is periodic, barrier of limited thickness - solution does not continue decaying to zero. There is tunneling between wells.

$$\psi = C \sinh(\gamma x) + D \cosh(\gamma x) \quad \frac{d\psi}{dx} = C\gamma \cosh(\gamma x) + D\gamma \sinh(\gamma x)$$

$$\gamma = \frac{\sqrt{2m(V_0 - E)}}{\hbar}$$

ψ must be continuous at $x = 0$, so $B = D$.

Also, $\frac{d\psi}{dx}$ must be continuous at $x = 0$, so $A\alpha = C\gamma$ or $C = (\alpha/\gamma)A$

From Bloch's theorem (Periodic potential)

$$\psi(x) = U(x)e^{ikx}$$

and

$$\psi(a) = e^{jk(a+b)}\psi(-b)$$

$$\left. \frac{d\psi}{dx} \right|_a = e^{jk(a+b)} \left. \frac{d\psi}{dx} \right|_{-b}$$

Therefore

$$A \sin \alpha a + B \cos \alpha a = e^{jk(a+b)} (A \sinh(-\gamma b) + B \cosh(-\gamma b))$$

$$A \left[\sin(\alpha c) + \frac{\alpha}{\gamma} e^{jk(a+b)} \sinh(\gamma b) \right] + B \left[\cos(\alpha c) - e^{jk(a+b)} \cosh(\gamma b) \right] = 0 \quad (42)$$


and

$$A\alpha [\cos(\alpha c) - e^{jk(a+b)} \cosh(\gamma b)] + B[-\alpha \sin(\alpha c) + \gamma e^{jk(a+b)} \sinh(\gamma b)] = 0 \quad (43)$$

Equations (42) and (43) have a non-trivial solution (i.e. a solution other than $A=B=0$ only if

$$\frac{\gamma^2 - \alpha^2}{2\alpha\gamma} \sin(\alpha c) \sinh(\gamma b) + \cos(\alpha c) \cosh(\gamma b) = \cos(k(b+c)) \quad (44)$$

At this point it is convenient to consider the special case in which $b \rightarrow 0$ and $V_0 \rightarrow \infty$ (makes things a little simpler – no problem for $E > V_0$ transition) while the product $V_0 b$ remains constant.




Letting $P = (mV_0bc)/\hbar^2$, (44) becomes

$$\frac{P}{\alpha c} \sin(\alpha c) + \cos(\alpha c) = \cos(kc)$$

(term 1)

(term 2)

This can be solved graphically. Solutions for $P = 3\pi/2$ (corresponding to a high barrier) are shown below. We plot (1) and (2) as function of αc .




We can only have a Solution for this when

$$\frac{P}{\alpha c} \sin(\alpha c) + \cos(\alpha c) \leq 1$$

and as $\cos(kc) \leq 1$ always we have a valid solution for ranges of αc which implies a valid solution for particular energy regions.

E




We therefore find there are α 's for which there is no valid K. As $E = \frac{\hbar^2}{2m} \alpha^2$

This means there are disallowed regions of energy, i.e., energy gaps.

Energies and wave functions for electrons in crystal (periodic potential $V(x)$)

a) Isolated Potential well



- only discrete energies E allowed
- wave functions are standing waves

b) Free Electrons

$$\psi(x) = Ce^{jkx}$$


$$\Psi(x,t) = Ce^{j(kx - \omega t)}$$

- plane wave solutions


- any energy allowed:

$$E = \frac{\hbar^2 k^2}{2m}$$

- Parabolic E versus k



c) Periodic Potential




- ability of electrons to tunnel between barrier walls spreads out the discrete energy levels seen for isolated wells into bands

- We find a number of bands of energies are allowed.
- We have restricted k values: $(-\pi/a \leq k \leq \pi/a)$
- The wave functions are Bloch waves: $\psi_k(x) = U_k(x)Ce^{ikx}$

$\psi_k(x, t) = U_k(x)e^{j(kx - \omega t)}$ - which is modulated travelling wave

- Wave functions act like free electrons (almost)

Simplified energy diagram:



General results from Kronig-Penney model:

- if potential barrier between wells is strong, energy bands are narrowed and spaced far apart

(Corresponds to crystals in which electrons are tightly bound to ion cores, and wavefunctions do not overlap much with adjacent cores. Also true for lowest energy bands)

- if potential barrier between wells is weak, energy bands are wide and spaced close together (this is typically situation for metals with weakly bound electrons – e.g. alkali metals. Here the “nearly free” electron model works well.)